

AD-A077 138

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

GLOBAL STABILITY OF STATIONARY SOLUTIONS OF REACTION-DIFFUSION --ETC(U)

SEP 79 R A GARDNER

DAAG29-75-C-0024

UNCLASSIFIED

MRC-TSR-1993

NL

| OF |  
AD  
A077138



END  
DATE  
FILMED  
12-79  
DDC

AD A 077138

MRC Technical Summary Report #1993 ✓

GLOBAL STABILITY OF STATIONARY SOLUTIONS  
OF REACTION-DIFFUSION SYSTEMS

Robert A. Gardner

Mathematics Research Center  
University of Wisconsin-Madison ✓  
610 Walnut Street  
Madison, Wisconsin 53706

September 1979

(Received June 20, 1979)



Approved for public release  
Distribution unlimited

Sponsored by

U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709

National Science Foundation  
Washington, D.C. 20550

DDC FILE COPY

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

GLOBAL STABILITY OF STATIONARY SOLUTIONS  
OF REACTION-DIFFUSION SYSTEMS

Robert A. Gardner<sup>\*</sup>

Technical Summary Report #1993  
September 1979

ABSTRACT

We give a condition <sup>is given</sup> which implies that the trivial solution,  $U \equiv 0$ , <sup>is identically equal to</sup> of a class of reaction-diffusion systems with homogeneous Dirichlet boundary conditions, is a global attractor for all non-negative solutions. In certain cases, this condition, which relates the diffusion matrix and the domain to a parameter which depends on the nonlinear term, significantly improves similar conditions which can be obtained from energy estimates. Applications are given to equations arising in mathematical ecology.

AMS (MOS) Subject Classifications - 35B30, 35B40, 35K50, 35K55

Key Words - Reaction-diffusion systems, stability, boundary values

Work Unit Number 1 - Applied Analysis

<sup>\*</sup> Department of Mathematics, Heriot-Watt University, Riccarton, Currie, Edinburgh EH14 4AS, Scotland.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024. This material is based upon work supported by the National Science Foundation under Grant No. MCS78-09525.

## SIGNIFICANCE AND EXPLANATION

Reaction-diffusion systems are systems of nonlinear partial differential equations which arise in various aspects of science and engineering, including mathematical ecology. Such equations can be used to describe the evolution of interacting and diffusing species in a bounded domain, together with the usual assumption that there is no movement across the boundary; however, when there is migration across the boundary, other boundary conditions are more appropriate. It is therefore of interest to study the behavior of solutions of such equations with a variety of different boundary conditions; that is, we suppose that the density of each species at the boundary is zero.

The purpose of this paper is to give a condition which implies that each species in the system moves to extinction. Such conditions are easily obtained from standard "energy" estimates and have been observed by a number of authors. We shall give a condition which, for a reasonable class of equations, provides significantly sharper results. In particular, under certain conditions we obtain the global stability of extinction for any domain and for arbitrary diffusion rates. Applications are given to equations which describe the ecological interactions of predation and competition.

Accession For	
NTIS	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Availand/or special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.



GLOBAL STABILITY OF STATIONARY SOLUTIONS  
OF REACTION-DIFFUSION SYSTEMS

Robert A. Gardner\*

I.

The purpose of this paper is to give a condition which implies that the trivial solution  $U \equiv 0$  of the reaction-diffusion system

$$U_t = D\Delta U + F(U), \quad U(0, x) = U^0(x), \quad (1)$$

$$U|_{\partial\Omega \times \mathbb{R}_+} = 0, \quad (2)$$

is a global attractor for all non-negative solutions of (1), (2). Our condition relates the diffusion matrix,  $D$ , and  $\Omega$  to a third parameter which depends on the nonlinear term. In certain cases, this provides a considerable improvement over the conditions and results obtained from energy estimates by Conway, Hoff and Smoller [3]. These results have applications to systems arising in mathematical ecology.

We assume that  $D = \text{diag}(d_1, \dots, d_n)$ , where each  $d_i$  is a positive constant,  $F(U) = (f_1(U), \dots, f_n(U))$ ,  $\Delta$  is the  $m$ -dimensional Laplacian, and that  $\Omega$  is a bounded domain in  $\mathbb{R}^m$  lying on one side of its boundary, which we assume to be smooth. The interactive term  $F$  is assumed to be smooth and to possess bounded invariant sets  $\Sigma$  of the form  $\Sigma = \prod_{i=1}^n [0, a_i]$ , for sufficiently large  $a_i > 0$ ; that is, if  $U^0(x)$  is continuous and has values in  $\Sigma$  then  $U(x, t) \in \Sigma$  for  $x \in \Omega$  and all  $t \geq 0$  for which the solution  $U$  of (1), (2) exists. This will be the case if we impose the condition

$$F(U) \cdot \nu(U) \leq 0, \quad (3a)$$

for all  $U \in \partial\Sigma$  and  $x \in \Omega$ ; here,  $\nu(U)$  is an outward normal to  $\partial\Sigma$  at  $U$ .

It follows that there then exists a unique, smooth solution of (1), (2) defined

---

\*Department of Mathematics, Heriot-Watt University, Riccarton, Currie, Edinburgh EH14 4AS, Scotland.

for all  $t \geq 0$  and with values in  $\Sigma$ ; see [2] for details. It will also be assumed that  $f_i$  has the form

$$f_i(U) = u_i M_i(U) \quad (3b)$$

In the context of mathematical ecology,  $M_i$  is the local growth rate of the  $i^{\text{th}}$  species.

We first give a condition which implies that the plane  $\{u_i = 0\}$  is a global attractor for solutions of (1), (2) with data  $U^0(x) \in \Sigma$ . Let

$$g_i(w) = \sup\{f_i(\xi_1, \dots, \xi_{i-1}, w, \xi_{i+1}, \dots, \xi_n) : 0 \leq \xi_j \leq a_j, i \neq j\}, \quad (4)$$

and define  $G_i(w) = \int_0^w g_i(s) ds$ . Now let

$$\lambda_i = \sup\{G_i(w)/w^2 : 0 \leq w \leq a_i\}.$$

(Note that  $\lambda_i$  is bounded by an expression which depends on  $a_i$  and the Lipschitz constant of  $g_i$ , which is finite; see [4, Lemma].) If  $R$  is the radius of the smallest ball containing  $\Omega$ , our condition is that

$$\lambda_i R^{2-1/\gamma} < 1, \quad (5)$$

where  $\gamma > 0$  depends only on  $m$  if  $m > 1$ ; (if  $m = 1$   $\gamma$  also depends on  $g_i$  and  $R^{2-1/\gamma}$ ). The proof employs a comparison technique introduced by Conway and Smoller, [4], to estimate  $u_i$  from above by the solution  $w$  of the scalar equation

$$w_t = d_i \Delta w + g_i(w), \quad w(x, 0) = u_i^0(x), \quad w|_{\partial\Omega \times \mathbb{R}_+} = 0. \quad (6)$$

Next, it is shown that condition (5) implies that zero is the unique solution of the steady state equation

$$0 = d_i \Delta w + g_i(w), \quad w|_{\partial\Omega} = 0, \quad (7)$$

associated with (6), and hence, that  $w(x, t)$  must decay to zero as  $t$  approaches infinity. Thus if (5) holds for  $1 \leq i \leq n$ , the origin must be a global attractor. It should be noted that if  $\lambda_i \leq 0$ , (which occurs in a number of

examples), our condition is independent of  $D$  and  $\Omega$ . In Section III, we apply the above result to equations which describe the ecological interactions of predation and competition.

Acknowledgement. The author would like to thank M. Crandall for suggesting that  $D$  and  $\Omega$  be related to  $\lambda_1$ ; (the theorem was originally proved with the hypothesis  $\lambda_1 \leq 0$  instead of (5)).

## II.

We shall begin with a discussion of the case  $m = 1$  and  $\lambda_1 \leq 0$ . The proof, though trivial, provides a simple geometric interpretation of condition (5) which may help to extend our results to a more general setting. The crucial step is to show that zero is the unique non-negative solution of (7). If  $w$  is a smooth non-negative solution on  $\Omega = (-l, l)$  we multiply (7) by  $w_x$  and integrate from zero to  $x$  to obtain

$$C = d_i w_x^2/2 + G_i(w)$$

where  $C = d_i w_x^2(0)/2 + G_i(w(0))$ . Hence the solutions of our boundary value problem coincide with the level curves of the expression  $d_i z^2/2 + G_i(w)$ , where  $z = w_x$ . When  $G_i(w) \leq 0$ , the level curves of this expression are as in Figure 1;




Figure 1.



the solutions of (7) must cross the  $z$ -axis when  $x = \pm l$ . Clearly, there is no such smooth non-negative solution. We shall prove a similar result in several space variables by reducing the problem to one dimension.

Theorem. Suppose that conditions (3) and (5) hold. Then  $\lim_{t \rightarrow \infty} u_i(x, t) = 0$ , uniformly for  $x \in \Omega$ , where  $u_i$  is the  $i^{\text{th}}$  component of the solution of (1), (2).

Proof. We begin by noting that (3) together with the definition (4) of  $g_i$  imply that  $g_i(0) = 0$  and that  $g_i(a_i) \leq 0$ , so that 0 and  $a_i$  are respectively lower and upper solutions of (7). Moreover, it is shown in [4] that  $g_i$  is uniformly Lipschitz (and hence Hölder) continuous on  $[0, a_i]$ , so that there exists a unique classical solution  $w$  of (6) with values in  $[0, a_i]$  defined for all  $t \geq 0$ .

The desired comparison  $u_i \leq w$  is obtained by arguing as in [4]. In particular, we let  $z = u_i - w$  and  $h(x, t, z) = f_i(u) - g_i(w)$ , so that  $z$  satisfies the equation

$$z_t = d_i \Delta z + h(x, t, z), \quad z|_{\partial\Omega \times \mathbb{R}_+} = z|_{t=0} = 0. \quad (8)$$

From (4), we see that  $h(x, t, 0) \leq 0$ , so that  $\{z \leq 0\}$  is an invariant set for (8); hence  $u_i < w$  in  $\Omega \times \mathbb{R}_+$ .

We now assert that zero is the unique smooth solution of

$$-d_i \Delta w = g_i(w), \quad w|_{\partial\Omega} = 0, \quad (9)$$

with values in  $[0, a_i]$ . Suppose that this is not the case, and let  $w_0(x) \not\equiv 0$  be such a solution. Let  $D = \{x \in \mathbb{R}^m : |x| < R\}$ , where  $R > 0$  is chosen such that  $\Omega \subset D$ , and define

$$w_*(x) = \begin{cases} w_0(x), & x \in \Omega \\ 0, & x \in D \setminus \Omega. \end{cases}$$

Clearly,  $w_* \in H_0^1(D)$ . We claim that  $w_*$  is a weak lower solution of

$$-d_i \Delta w = g_i(w), \quad w|_{\partial D} = 0; \quad (9)_D$$



that is,  $w_*$  satisfies the inequality

$$\int_D d_i \nabla w_* \cdot \nabla \varphi dx \leq \int_D g_i(w_*) \varphi dx ,$$

for all  $\varphi \in H_0^1(D)$  with  $\varphi \geq 0$ . It clearly suffices to prove this inequality for all  $\varphi \in C_0^\infty(D)$ . Let  $n$  denote the outward unit normal to  $\partial\Omega$ . Then

$dw_0/dn \leq 0$ , since  $w_0 \geq 0$  in  $\Omega$  and  $w_0 = 0$  on  $\partial\Omega$ . Let  $K = \text{supp } \varphi$ , and let  $H = \partial\Omega \cap K$ . If we give  $H$  the orientation induced by  $n$ , we have that

$$\int_{\partial(\Omega \cap K)} d_i \varphi dw_0/dn d\sigma = \int_H d_i \varphi dw_0/dn d\sigma \leq 0 ,$$

since  $\varphi = 0$  on  $\partial K$ . Thus

$$\begin{aligned} \int_D d_i \nabla \varphi \cdot \nabla w_* dx &= \int_{K \cap \Omega} d_i \nabla \varphi \cdot \nabla w_* dx + \int_{K \setminus \Omega} d_i \nabla \varphi \cdot \nabla w_* dx \\ &= \int_{\partial(K \cap \Omega)} d_i \varphi dw_0/dn d\sigma - \int_{K \cap \Omega} d_i \varphi \Delta w_0 dx \\ &\leq \int_{K \cap \Omega} d_i \varphi g_i(w_0) dx = \int_D d_i \varphi g_i(w_*) dx , \end{aligned}$$

since  $g_i(0) = 0$ . Thus  $w_*$  is an  $H_0^1(D)$  lower solution of  $(9)_D$ . Clearly  $w^* = a_i$  is an upper solution, so that we may apply an existence theorem of Dueul and Hess, [7], to obtain an exact  $H_0^1(D)$  solution  $w_1$  of  $(9)_D$  which satisfies  $w_* \leq w_1 \leq w^*$ . Since  $w_1$  is bounded and  $g_i$  is Lipschitz continuous, the usual bootstrapping arguments imply that  $w_1 \in C^{2+\alpha}(D)$ , so that by an existence theorem of Amann, [1], there exists a maximal  $C^{2+\alpha}(D)$  solution  $w$  satisfying  $w_1 \leq w \leq w^*$ . Since any rotation of a solution of  $(9)_D$  is again a solution of  $(9)_D$ , it follows that  $w$  is rotationally invariant, so that  $w$  satisfies the O.D.E.

$$-d_i(w_{rr} + \frac{m-1}{r} w_r) = g_i(w), \quad w(R) = 0, \quad w_r(0) = 0 ;$$

(the boundary condition at  $r = 0$  follows from the smoothness of  $w$ ). We rescale this equation to obtain

$$-p(w_{rr} + \frac{m-1}{r} w_r) = g_i(w), \quad w(1) = 0, \quad w_r(0) = 0 , \quad (10)$$

where  $p = d_i R^{-2}$ . Now multiply (10) by  $w_r$  and integrate from  $r = 0$  to  $r = 1$  to obtain

$$p(w_r^2(1)/2 + (m-1) \int_0^1 s^{-1} w_r^2(s) ds) = G_i(w(0)). \quad (11)$$

(We have used the fact that  $G_i(w(1)) = G_i(0) = 0$ ). First, we shall suppose that  $m > 1$ . Since  $w(1) = 0$ , by Poincaré's inequality there exists a constant

$C_1 > 0$  such that  $C_1 \int_0^1 w^2(s) ds \leq \int_0^1 w_r^2(s) ds$ , so that

$$C_1(1 + C_1)^{-1} \int_0^1 (w^2 + w_r^2) ds \leq \int_0^1 w_r^2(s) ds \leq \int_0^1 s^{-1} w_r^2(s) ds.$$

By the Sobolev embedding theorem, there exists a constant  $C_2 > 0$  such that

$C_2 \|w\|_\infty \leq \|w\|_{H^1}$ . The last two inequalities together with (11) imply that

$p\gamma^{-1}w(0)^2 \leq G_i(w(0))$ , where  $\gamma^{-1} = C_1 C_2^2(m-1)/(C_1 + 1)$ ; note that  $\gamma$  depends

only on  $m$ . Suppose that  $w(0) = 0$ . Then there exists  $r_m$  with  $0 < r_m < 1$

and such that  $\|w\|_\infty = w(r_m)$ . Multiply (10) by  $w_r$  and integrate from  $r = 0$  to  $r = r_m$  to obtain

$$-p(m-1) \int_0^{r_m} s^{-1} w_r^2(s) ds = G_i(w(r_m)) < 0,$$

and from  $r = r_m$  to  $r = 1$  to obtain

$$p(w_r^2(1)/s + (m-1) \int_{r_m}^1 s^{-1} w_r^2(s) ds) = G_i(w(r_m)) > 0,$$

contradiction. Thus we have that  $w(0) > 0$  and that

$$1 \leq p^{-1} \gamma G_i(w(0)) w(0)^{-2} \leq R^2 \gamma \lambda_i / d_i.$$

This inequality violates condition (5), yielding the desired contradiction.

If  $m = 1$ , we must use a different argument. Let  $F(w, z) = (z, -p^{-1}g_i(w))$ , so that the solution of (10) also satisfies

$$(w, z)_r = F(w, z), \quad (w(1, \beta), z(1, \beta)) = (0, \beta), \quad (12)$$

when  $\beta = \beta_0$  for some  $\beta_0 < 0$ . Since solutions to (12) are unique, solutions of (12) are uniformly bounded by the particular solution of (12) when  $\beta = \beta_0$  for any  $\beta$  with  $0 \geq \beta \geq \beta_0$ . Viewing  $\beta$  as parameter, we apply Gronwall's inequality to obtain for any  $\beta_1, \beta_2$  between zero and  $\beta_0$

$$(w(0, \beta_1) - w(0, \beta_2))^2 + (z(0, \beta_1) - z(0, \beta_2))^2 \leq \gamma |\beta_1 - \beta_2|^2,$$

where  $\gamma$  depends only the Lipschitz constant of  $p^{-1}g_i$  and on  $a_i$ . Taking  $\beta_2 = 0$  and  $\beta_1 = \beta_0$ , we have that

$$\gamma^{-1} w(0)^2 \leq w_r(1)^2.$$

Since  $m = 1$ , we clearly have that  $w(0) \neq 0$ . The proof now proceeds as above.

We shall now show that  $\lim_{t \rightarrow \infty} w(x, t) = 0$ , pointwise, where  $w(x, t)$  is the solution of (6). If this is not the case, there exists  $x_0 \in \Omega$  such that  $\limsup_{t \rightarrow \infty} w(x_0, t) > 0$ . By [8, Lemma 3.8], there exists a constant  $K > 0$  such that  $|\nabla_x w(x, t)| < K$  for  $(x, t) \in \Omega \times [1, \infty)$ . We may therefore choose a sequence  $\{t_n\}$  with  $\lim_{n \rightarrow \infty} t_n = \infty$ , and such that if  $w_n(x) = w(x, t_n)$ , then  $\lim_{n \rightarrow \infty} w_n(x) = w_0(x)$  in  $C^\alpha(\Omega)$ ,  $0 < \alpha < 1$ , and  $w_0(x_0) > 0$ . We claim that  $w_0(x)$  is a solution of (9). If  $\varphi \in C_0^\infty(\Omega)$ , then

$$\begin{aligned} \left| \int_{\Omega} [d_i \Delta \varphi w_0 + \varphi g_i(w_0)] dx \right| &= \lim_n \left| \int_{\Omega} [d_i \Delta \varphi w_n + \varphi g_i(w_n)] dx \right| \\ &= \lim_n \left| \int_{\Omega} d_i \varphi w_t(x, t_n) dx \right| \\ &\leq \lim_n d_i \|w_t(\cdot, t_n)\|_{L^2} \|\varphi\|_{L^2}. \end{aligned}$$

Now multiply (6) by  $w_t$ , integrate over  $\Omega \times [1, \infty)$ , and apply Green's theorem to obtain

$$\int_1^\infty \int_{\Omega} w_t^2 dx dt = \lim_{T \rightarrow \infty} \left[ \int_{\Omega} (d_i |\nabla w|^2 / 2 + G_i(w)) dx \right]_{t=1}^{t=T} < \infty,$$

so that  $\lim_{t \rightarrow \infty} \|w_t(\cdot, t)\|_{L^2} = 0$ . Hence  $w_0$  is a weak (and therefore strong)



solution of (9); since  $g_i$  is Lipschitz continuous and  $w_0$  is Hölder continuous, we must therefore have that  $w_0$  is a non-trivial, non-negative classical solution, yielding the desired contradiction.

We obtain uniform decay of  $w$  as follows. If  $\mu(t) = \|w(\cdot, t)\|_{L^2}^2$ , we obtain the differential inequality  $\mu' + C\mu \leq \int_{\Omega} w g_i(w) dx$  by multiplying (6) by  $w$ , integrating over  $\Omega$ , integrating by parts, and by finally applying Poincaré's inequality. Since  $w g_i(w)$  is a bounded function which converges pointwise to zero, the Lebesgue dominated convergence theorem implies that  $\lim_{t \rightarrow \infty} \int_{\Omega} w g_i(w) dx = 0$ , and thus the above differential inequality implies that  $\lim_{t \rightarrow \infty} \mu(t) = 0$ . Since  $w$  is bounded, we obtain  $L^p$  decay for any  $p \geq 2$ , and we obtain uniform decay from [9, Lemma 3.1].  $\square$

Our result can be extended in several ways. The functions  $f_i(U)$  may be allowed to depend on  $x$  and  $t$ ; we need only alter the definition (4) of  $g_i$  by taking the supremum of  $f_i(U, x, t)$  over all appropriate values of  $x$  and  $t$  in addition to the variables  $u_j$ ,  $j \neq i$ .

We may also replace  $-d_i \Delta$  by an elliptic operator  $L_i$  of the form

$$L_i = - \sum_{j,k} a_{j,k}^i \partial^2 / \partial x_i \partial x_j,$$

where the  $a_{j,k}^i$ 's are constants which satisfy the condition  $\sum_{j,k} a_{j,k}^i \xi_j \xi_k \geq d_i |\xi|^2$ . We modify the proof by first performing a linear change of variables  $x \rightarrow x'$  which transforms  $L_i$  into  $-d_i \Delta$ .

We can also consider other boundary value problems when  $m = 1$ ; (2) is replaced by the mixed condition

$$u_i + \beta_i(x) du_i / dn = 0, \quad x = \pm l,$$

where  $\Omega = (-l, l)$  and  $\beta_i \geq 0$ . For simplicity, we consider the case  $\lambda_i \leq 0$ . Suppose that the condition

$$\|\beta_i\|_{\infty} < \max\{|g_i(w)(-G_i(w))^{-1/2}| : 0 \leq w \leq a_i\} \quad (13)$$

holds. We claim that there are no non-negative, non-trivial stationary solutions. Such a solution must connect the ray  $w = \beta_i(-l)z$  when  $x = -l$  to the ray  $w = -\beta_i(l)z$  when  $x = l$ , where  $w_x = z$ . Condition (13) ensures that the former ray lies above those trajectories which coincide with the curve OAB in Figure 1 and that the latter ray lies below those trajectories which coincide with the curve OCD in Figure 1, since the slope of OAB and OCD is  $\pm g_i(w)(-G_i(w))^{-1/2}$  when  $G_i(w) \neq 0$ , and the appropriate right or left hand limit of this expression when  $G_i(w) = 0$ . It should be noted that condition (13) is independent of  $D$  and  $\Omega$ .

The above remarks indicate that our theorem should be true in somewhat greater generality. For example, it is reasonable to consider uniformly elliptic variable coefficient operators of the form  $L_i = -\sum_{j,k} a_{j,k}^i(x,t) \partial^2 / \partial x_i \partial x_j$ , or mixed boundary conditions of the form (13) in several space variables. However, the proof given above doesn't seem to be adaptable to such situations. It seems likely that a proof might be found which avoids using the extension procedure and the rotational symmetry of the Laplacian.

### III.

In this section, we shall consider a few examples arising in mathematical ecology. Most attention has been given to the Cauchy problem and to the initial-boundary value problem with homogeneous Neumann, (or "no flux") conditions, [4], [5], [6]. However, Dirichlet or mixed boundary conditions may be more appropriate if there is migration across the boundary. It is worth noting that our result is not valid for the pure Neumann or Cauchy problem, since in these cases, there may exist non-trivial constant rest states.

#### EXAMPLE 1. PREDATION.

Let  $v$  be the population density of a predator species, and let  $u$  be the density of its prey. Let  $M$  and  $N$  be the growth rates of  $u$  and  $v$  respectively; then  $u$  and  $v$  satisfy the system

$$\begin{aligned} u_t &= d_1 \Delta u + uM(u,v) \\ v_t &= d_2 \Delta v + vN(u,v) \end{aligned} \quad (14)$$

The interaction is characterized by making certain assumptions about the algebraic signs of  $M$  and  $N$  and their partial derivatives; for a complete discussion, see [4], [5], [6]. We shall assume that (i),  $M \leq 0$  and  $N \leq 0$  if  $u$  and  $v$  are both near zero, (ii),  $M \leq 0$  if  $u$  is large and  $N \leq 0$  if  $v$  is large, and (iii),  $M_v \leq 0$  and  $N_u \geq 0$ . Condition (i) requires that there be a critical population density below which each species goes extinct and (ii) is a resource limitation condition. The predator prey relationship is characterized by (iii). One such example is the Rosensweig-MacArthur equations, where we take

$$M = u(-v + \delta(u - a)(b - u))$$

$$N = v(-v + cu - d) ;$$

$a, b, c, d$ , and  $\delta$  are positive constants. The phase diagram of the vector field  $(uM, vN)$  is given in Figure 2.




Figure 2.



We have that  $g_1(w) = \delta w(w - a)(b - w)$ . A simple computation shows that  $\lambda_1 = \delta[(a + b)^2/9 - ab/2]$ , so that  $\lambda_1 \leq 0$  if  $a \geq b/2$ . In this case,  $u$  must tend to zero independently of  $D, \Omega$ , and  $\delta$ . Since  $N \leq 0$  if  $u$  is sufficiently small, we must have that  $v$  tends to zero also, (even though  $\lambda_2$  is always positive here).

#### EXAMPLE 2. COMPETITION.

We now let  $u$  and  $v$  be the population densities of two competing species. These variables will again satisfy a system of equations of the form (14). However, we now replace hypothesis (iii) with (iii)',  $M_v \leq 0$  and  $N_u \leq 0$ . (We still assume that (i) and (ii) hold). The phase diagram of an example of such a system is indicated in Figure 3.




Figure 3.

For definiteness, let

$$M = u(-v + \delta_1(u - a_1)(b_1 - u))$$

$$N = v(-u + \delta_2(v - a_2)(b_2 - v)) .$$

As in Example 1, we have that the origin is a global attractor, provided that  $a_i \geq b_i/2$ ,  $i = 1, 2$ .

# REFERENCES

- [1] H. Amann, On the existence of positive solutions of nonlinear elliptic boundary value problems, Indiana Univ. Math. J., 21 (1971), 125-146.
- [2] K. N. Chueh, C. C. Conley, and J. A. Smoller, Positively invariant regions for systems of nonlinear diffusion equations, Indiana Univ. Math. J., 26 (1977), 373-391.
- [3] E. Conway, D. Hoff, and J. A. Smoller, Large time behavior of solutions of systems of nonlinear reaction-diffusion equations, SIAM J. Appl. Math., 35 (1978), 1-16.
- [4] E. Conway and J. A. Smoller, A comparison technique for systems of reaction-diffusion equations, Comm. Partial Differential Equations, 2 (1977), 679-697.
- [5] \_\_\_\_\_, Diffusion and the classical ecological interactions, Proceedings of Conference on Nonlinear Diffusion, Pitman, London.
- [6] \_\_\_\_\_, Diffusion and the predator-prey interaction, SIAM J. Appl. Math., 33 (1977), 673-686.
- [7] J. Dueul and P. Hess, A criterion for the existence of solutions of nonlinear elliptic boundary value problems, Proceedings of the Royal Society of Edinburgh, 74A, 3, 1974/75, 49-54.
- [8] R. A. Gardner, Asymptotic behavior of systems of nonlinear reaction-diffusion equations with Dirichlet conditions, Ph.D. Dissertation, University of Michigan, 1978.
- [9] C. Kahane, On a system of nonlinear parabolic equations arising in chemical engineering, J. Math. Anal. Appl., 53 (1976), 343-358.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1993	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) GLOBAL STABILITY OF STATIONARY SOLUTIONS OF REACTION-DIFFUSION SYSTEMS.		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) 10 Robert A. Gardner		8. CONTRACT OR GRANT NUMBER(s) 15 DAAG29-75-C-0024, NSF-MCS78-09525
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18.		12. REPORT DATE 11 September 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9 Technical Summary Rept,		13. NUMBER OF PAGES 12
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 14 MRC-TSR-1993		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE 12 16
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U.S. Army Research Office National Science Foundation P.O. Box 12211 Washington, D.C. 20550 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Reaction-diffusion systems, stability, boundary values		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We give a condition which implies that the trivial solution, $U \equiv 0$ , of a class of reaction-diffusion systems with homogeneous Dirichlet boundary conditions, is a global attractor for all non-negative solutions. In certain cases, this condition, which relates the diffusion matrix and the domain to a parameter which depends on the nonlinear term, significantly improves similar conditions which can be obtained from energy estimates. Applications are given to equations arising in mathematical ecology.		